

Centers for Disease Control and Prevention

April 13, 2020

Return to Work Criteria for HCP with Confirmed or Suspected COVID-19

Use the *Test-based strategy* as the preferred method for determining when HCP may return to work in healthcare settings:

1. *Test-based strategy*. Exclude from work until
 - Resolution of fever without the use of fever-reducing medications **and**
 - Improvement in respiratory symptoms (e.g., cough, shortness of breath), **and**
 - Negative results of an FDA Emergency Use Authorized molecular assay for COVID-19 from at least two consecutive nasopharyngeal swab specimens collected ≥ 24 hours apart (total of two negative specimens) [1]. See [Interim Guidelines for Collecting, Handling, and Testing Clinical Specimens for 2019 Novel Coronavirus \(2019-nCoV\)](#).

If the *Test-based strategy* cannot be used, the *Non-test-based strategy* may be used for determining when HCP may return to work in healthcare settings:

2. *Non-test-based strategy*. Exclude from work until
 - At least 3 days (72 hours) have passed *since recovery* defined as resolution of fever without the use of fever-reducing medications **and** improvement in respiratory symptoms (e.g., cough, shortness of breath); **and**,
 - At least 7 days have passed *since symptoms first appeared*

HCP with laboratory-confirmed COVID-19 who have not had any symptoms should be excluded from work until 10 days have passed since the date of their first positive COVID-19 diagnostic test assuming they have not subsequently developed symptoms since their positive test.

If HCP had COVID-19 ruled out and have an alternate diagnosis (e.g., tested positive for influenza), criteria for return to work should be based on that diagnosis.

Return to Work Practices and Work Restrictions

After returning to work, HCP should:

- Wear a facemask for source control at all times while in the healthcare facility until all symptoms are completely resolved or until 14 days after illness onset, whichever is longer. A facemask instead of a cloth face covering should be used by these HCP for source control during this time period while in the facility. After this time period, these HCP should revert to their facility policy regarding [universal source control](#) during the pandemic.
 - A facemask for source control does not replace the need to wear an N95 or higher-level respirator (or other recommended PPE) when indicated, including when caring for patients with suspected or confirmed COVID-19.
 - Of note, N95 or other respirators with an exhaust valve might not provide source control.
- Be restricted from contact with severely immunocompromised patients (e.g., transplant, hematology-oncology) until 14 days after illness onset.

- Self-monitor for symptoms and seek re-evaluation from occupational health if respiratory symptoms recur or worsen.

Strategies to Mitigate Healthcare Personnel Staffing Shortages

Maintaining appropriate staffing in healthcare facilities is essential to providing a safe work environment for healthcare personnel (HCP) and safe patient care. As the COVID-19 pandemic progresses, staffing shortages will likely occur due to HCP exposures, illness, or need to care for family members at home. Healthcare facilities must be prepared for potential staffing shortages and have plans and processes in place to mitigate them, including considerations for permitting HCP to return to work without meeting all return to work criteria above. Refer to the *Strategies to Mitigate Healthcare Personnel Staffing Shortages* document for information.

Footnotes

¹All test results should be final before isolation is ended. Testing guidance is based upon limited information and is subject to change as more information becomes available. In persons with a persistent productive cough, SARS-CoV-2-RNA might be detected for longer periods in sputum specimens than in upper respiratory tract (nasopharyngeal swab) specimens.

Definitions

Cloth face covering: Textile (cloth) cover that are intended to keep the person wearing one from spreading respiratory secretions when talking, sneezing, or coughing. **They are not PPE and it is uncertain whether cloth face coverings protect the wearer.** Guidance on design, use, and maintenance of cloth face coverings is available.

Facemask: Facemasks are PPE and are often referred to as surgical masks or procedure masks. Use facemasks according to product labeling and local, state, and federal requirements. FDA-cleared surgical masks are designed to protect against splashes and sprays and are prioritized for use when such exposures are anticipated, including surgical procedures. Facemasks that are not regulated by FDA, such as some procedure masks, which are typically used for isolation purposes, may not provide protection against splashes and sprays.

Respirator: A respirator is a personal protective device that is worn on the face, covers at least the nose and mouth, and is used to reduce the wearer's risk of inhaling hazardous airborne particles (including dust particles and infectious agents), gases, or vapors. Respirators are certified by the CDC/NIOSH, including those intended for use in healthcare.